

1

ORIGINAL FILE COPY

2

AD-A233 585

OFFICE OF NAVAL RESEARCH

Grant N00014-90-J-1193

TECHNICAL REPORT No. 43

Heisenberg Approach to Photon Emission Near a Phase Conjugator

by

H. F. Arnoldus and Thomas F. George

Prepared for publication

in


*Physical Review A*

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

March 1991

Reproduction in whole or in part is permitted for any purpose of the  
United States Government.

This document has been approved for public release and sale;  
its distribution is unlimited.



91 3 28 018

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/91/TR-43			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Grant N00014-90-J-1193		
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Heisenberg Approach to Photon Emission Near a Phase Conjugator					
12. PERSONAL AUTHOR(S) Henk F. Arnoldus and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) March 1991	
				15. PAGE COUNT 21	
16. SUPPLEMENTARY NOTATION Prepared for publication in <i>Physical Review A</i>					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	ATOM HEISENBERG APPROACH		
			PHASE CONJUGATOR PLANE-WAVE EXPANSION		
			PHOTON EMISSION CLASSICAL MAXWELL EQUATIONS		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) An expression for the emitted fluorescence radiation by an atom near a phase conjugator is derived from a plane-wave expansion of the electric field operator and Heisenberg's equation of motion for the annihilation operator. The result is compared to a solution which was found previously, based on the classical Maxwell equations, where it is shown that both theories yield the same expression for the field in the radiation zone, in the limit of a transparent medium, which confirms the correctness of either approach to the problem of optical phase conjugation of atomic radiation.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Physical Review A, in press

HEISENBERG APPROACH TO PHOTON EMISSION NEAR A PHASE CONJUGATOR

Henk F. Arnoldus  
Department of Physics  
Mendel Hall  
Villanova University  
Villanova, Pennsylvania 19085

and

Thomas F. George  
Departments of Chemistry and Physics & Astronomy  
239 Fronczak Hall  
State University of New York at Buffalo  
Buffalo, New York 14260

Accession For

ADONIS	<input checked="" type="checkbox"/>
PHYSICS	<input type="checkbox"/>
CHEMISTRY	<input type="checkbox"/>
ASTRONOMY	<input type="checkbox"/>

DATE	
BY	
REMARKS	

DATE

A-1

Abstract

An expression for the emitted fluorescence radiation by an atom near a phase conjugator is derived from a plane-wave expansion of the electric field operator and Heisenberg's equation of motion for the annihilation operator. The result is compared to a solution which was found previously, based on the classical Maxwell equations. It is shown that both theories yield the same expression for the field in the radiation zone, in the limit of a transparent medium. This confirms the correctness of either approach to the problem of optical phase conjugation of atomic radiation.

PACS: 42.65.H, 32.80, 52.50.D, 42.50.Kb

## I. INTRODUCTION

In a previous paper<sup>1</sup> we have studied the behavior of an atom near the surface of a four-wave mixing phase conjugator (PC). We calculated the fluorescent photon emission rate by solving Maxwell's equations for a dipole  $\underline{\mu}$  near a PC. The electric field  $\underline{E}(\underline{r}, t)$  was interpreted as a quantum operator field, with the argument that the classical Maxwell equations must be identical in form to the (quantum) Heisenberg equations of motion for the operator fields. Nevertheless, questions can be raised as to whether the identification of Maxwell's equations with Heisenberg's equations can be justified for the problem under consideration. In particular, the pump beams (with frequency  $\bar{\omega}$ ) for the four-wave mixing process are taken into account parametrically, and are represented by classical plane waves. This leads to factors of  $\exp(-2i\bar{\omega}t)$  in the expression for the fluorescence radiation field, rather than  $a(t)^2$ , with  $a(t)$  the annihilation operator for a photon in the pump beam. In addition, our results are in conflict with the results of Hendriks and Nienhuis,<sup>2,3</sup> who did not find the terms proportional to  $\underline{\mu}(t)^{(-)}$  (raising part) in the expression for the fluorescence field.

In this paper we solve the Heisenberg equation for the annihilation operator  $a_{\underline{k}\sigma}(t)$  for a photon with wave vector  $\underline{k}$  and polarization  $\sigma$  (either s = surface polarized or p = plane polarized). The solution is applied to evaluate the fluorescent radiation field in the far zone. These calculations are completely independent from our previous method, and the results can be used to verify the consistency of our approach.

## II. ELECTRIC FIELD

The electric-field operator  $\underline{E}(\underline{r}, t)$  can be represented as a sum of polarized plane waves. In the region  $z > 0$  (above the PC, where the atom is),

we have incident (inc) waves with wave vector  $\underline{k}$  and polarization  $\sigma$ . These waves give rise to specularly-reflected (r) waves, and to phase-conjugated (pc) waves which travel in the direction of  $-\underline{k}$ . In addition, there are transmitted (t) waves which have their origin in waves which are incident on the medium from the other side of the PC. The four-wave mixer produces also a nonlinear (nl) wave in  $z > 0$ .

The amplitude of each of the generated waves is related to the amplitude of the corresponding incident wave by a Fresnel coefficient. The electric field is explicitly,

$$\begin{aligned}
 \underline{E}(\underline{r}, t) = & \sum_{\underline{k}\sigma} \sqrt{\frac{\hbar\omega}{2\epsilon_0 V}} \left[ a_{\underline{k}\sigma}(t) \left\{ \underline{e}_{\underline{k}\sigma} e^{i\mathbf{k}\cdot\mathbf{r}} + R_{\underline{k}\sigma} \underline{e}_{-\underline{k}\sigma} e^{i\mathbf{k}\cdot\mathbf{r}} \right\} \right. \\
 & + a_{\underline{k}\sigma}^\dagger(t) e^{-2i\bar{\omega}t} P_{\underline{k}\sigma}^* \underline{e}_{-\underline{k}_{pc}\sigma} e^{-i\mathbf{k}_{pc}\cdot\mathbf{r}} \left. \right] \\
 & + \sum_{\underline{k}\sigma}'' \sqrt{\frac{\hbar\omega}{2\epsilon_0 V}} \left[ a_{\underline{k}\sigma}(t) T_{\underline{k}\sigma}' \underline{e}_{\underline{k}\sigma} e^{i\mathbf{k}\cdot\mathbf{r}} \right. \\
 & + a_{\underline{k}\sigma}^\dagger(t) e^{-2i\bar{\omega}t} N_{\underline{k}\sigma}'^* \underline{e}_{-\underline{k}_{nl}\sigma} e^{-i\mathbf{k}_{nl}\cdot\mathbf{r}} \left. \right] + \text{H.c.} \quad , \quad (2.1)
 \end{aligned}$$

where a prime (double prime) on the summation sign indicates a sum over waves which propagate in the  $z < 0$  ( $z > 0$ ) direction only. The unit polarization vectors  $\underline{e}_{\underline{k}_i\sigma}$  and the wave vectors  $\underline{k}_i$  are defined in Appendix A. The Heisenberg operator  $a_{\underline{k}\sigma}(t)$  is the annihilation operator for a photon which

is incident on the PC, either from  $z > 0$  or from the other side (back port) of the medium. We shall take the Schrödinger picture and Heisenberg picture to coincide at  $t = 0$ .

### III. EQUATION OF MOTION

The only unknown in expression (2.1) for  $\underline{E}(\underline{r}, t)$  is the annihilation operator  $a_{\underline{k}\sigma}(t)$  for  $t > 0$ . Its equation of motion is

$$i\hbar \frac{d}{dt} a_{\underline{k}\sigma}(t) = [a_{\underline{k}\sigma}(t), H] \quad , \quad (3.1)$$

with initial condition  $a_{\underline{k}\sigma}(0) = a_{\underline{k}\sigma}$ . The Hamiltonian  $H$  can be written as

$$H = H_a + H_r + H_{ar} \quad , \quad (3.2)$$

with

$$H_r = \sum_{\underline{k}\sigma} \hbar \omega_{\underline{k}\sigma} a_{\underline{k}\sigma}^\dagger a_{\underline{k}\sigma} \quad (3.3)$$

for the Hamiltonian of the radiation field, and where  $\omega = ck$ . The atomic Hamiltonian  $H_a$  can remain unspecified because it commutes with  $a_{\underline{k}\sigma}(t)$ . For the interaction, we take

$$H_{ar} = -\underline{\mu} \cdot \underline{E}(\underline{h}, 0) \quad , \quad (3.4)$$

in terms of the atomic dipole moment operator  $\underline{\mu}$  and the position  $\underline{h} = h\hat{e}_z$  of the atom. With Eqs. (3.3) and (3.4) the equation of motion becomes

$$i\hbar \frac{d}{dt} a_{\underline{k}\sigma}(t) = \hbar\omega a_{\underline{k}\sigma}(t) - \underline{\mu}(t) \cdot [\underline{a}_{\underline{k}\sigma}(t), \underline{E}(\underline{h}, t)] \quad , \quad (3.5)$$

where  $\underline{E}(\underline{h}, t)$  follows from Eq. (2.1) with  $\underline{r} = \underline{h}$ .

#### IV. SOLUTION

An integral of Eq. (3.5) is

$$a_{\underline{k}\sigma}(t) = e^{-i\omega t} \left( a_{\underline{k}\sigma} + \frac{i}{\hbar} \int_0^t dt' e^{i\omega t'} \underline{\mu}(t') \cdot [\underline{a}_{\underline{k}\sigma}(t'), \underline{E}(\underline{h}, t')] \right) \quad . \quad (4.1)$$

When we substitute Eq. (2.1) for  $\underline{E}(\underline{h}, t')$ , then it appears that many terms in the integrand oscillate at optical frequencies, and these terms average out to zero on a time scale of an optical cycle. The time dependence of the annihilation operator would be  $a_{\underline{k}\sigma}(t) = a_{\underline{k}\sigma} \exp(-i\omega t)$  in free evolution, which contains only a single positive frequency. When the interaction is taken into account,  $a_{\underline{k}\sigma}(t)$  acquires a spectral width around the frequency  $\omega$ , but it is still a positive-frequency operator. Similarly, when we split the dipole moment into a positive and a negative frequency part as

$$\underline{\mu}(t) = \underline{\mu}(t)^{(+)} + \underline{\mu}(t)^{(-)} \quad , \quad (4.2)$$

then  $\underline{\mu}(t)^{(+)} = \underline{\mu}^{(+)} \exp(-i\omega_0 t)$  in free evolution (for a two-level atom with transition frequency  $\omega_0$ ). The third time dependence enters as  $\exp(\pm 2i\bar{\omega}t)$ . Then we drop all terms in the integrand which oscillate with optical

frequencies, and retain the terms which oscillate with the difference of two optical frequencies. Furthermore, we notice that the first term on the right-hand side of Eq. (4.1) yields the vacuum field  $\underline{E}_v(\underline{r}, t)$ .

Therefore, we can write

$$\underline{E}(\underline{r}, t) = \underline{E}_v(\underline{r}, t) + \underline{E}_s(\underline{r}, t) \quad , \quad (4.3)$$

where  $\underline{E}_v(\underline{r}, t)$  is given by Eq. (2.1) with  $a_{\underline{k}\sigma}(t) \rightarrow a_{\underline{k}\sigma} \exp(-i\omega t)$ . The source field  $\underline{E}_s(\underline{r}, t)$  follows from the second term on the right-hand side of Eq. (4.1). We find explicitly,

$$\begin{aligned} \underline{E}_s(\underline{r}, t) = & \frac{i}{2\epsilon_0 V} \sum'_{\underline{k}\sigma} \omega e^{-i\omega t} b_{\underline{k}\sigma}(t) \left\{ \underline{e}_{\underline{k}\sigma} e^{i\underline{k} \cdot \underline{r}} + R_{\underline{k}\sigma} \underline{e}_{\underline{k}_r\sigma} e^{i\underline{k}_r \cdot \underline{r}} \right\} \\ & - \frac{i}{2\epsilon_0 V} \left( \sum'_{\underline{k}\sigma} \omega e^{-i(\omega-2\bar{\omega})t} b_{\underline{k}\sigma}(t) P_{\underline{k}\sigma} \underline{e}_{\underline{k}_{pc}\sigma} e^{i\underline{k}_{pc} \cdot \underline{r}} \right)^\dagger \\ & + \frac{i}{2\epsilon_0 V} \sum''_{\underline{k}\sigma} \omega e^{-i\omega t} b'_{\underline{k}\sigma}(t) T'_{\underline{k}\sigma} \underline{e}_{\underline{k}\sigma} e^{i\underline{k} \cdot \underline{r}} \\ & - \frac{i}{2\epsilon_0 V} \left( \sum''_{\underline{k}\sigma} \omega e^{-i(\omega-2\bar{\omega})t} b'_{\underline{k}\sigma}(t) N'_{\underline{k}\sigma} \underline{e}_{\underline{k}_{nl}\sigma} e^{i\underline{k}_{nl} \cdot \underline{r}} \right)^\dagger + \text{H.c.} \quad , \quad (4.4) \end{aligned}$$

in terms of the operators

$$b_{\underline{k}\sigma}(t) = \int_0^t dt' e^{i\omega t'} \underline{\mu}(t')^{(+)} \cdot \left( \underline{e}_{\underline{k}\sigma} e^{-i\underline{k} \cdot \underline{h}} + R_{\underline{k}\sigma}^* \underline{e}_{\underline{k}_r\sigma} e^{-i\underline{k}_r \cdot \underline{h}} \right)$$



$$+ \int_0^t dt' e^{i(\omega - 2\bar{\omega})t'} \underline{\mu}(t')^{(-)} \cdot (P_{\underline{k}\sigma}^* e_{\underline{k}_{pc}\sigma} e^{-i\mathbf{k}_{pc} \cdot \mathbf{h}}) \quad , \quad k_z < 0 \quad , \quad (4.5)$$

$$\begin{aligned} b'_{\underline{k}\sigma}(t) = & \int_0^t dt' e^{i\omega t'} \underline{\mu}(t')^{(+)} \cdot (T_{\underline{k}\sigma - \underline{k}\sigma}^* e^{-i\mathbf{k} \cdot \mathbf{h}}) \\ & + \int_0^t dt' e^{i(\omega - 2\bar{\omega})t'} \underline{\mu}(t')^{(-)} \cdot (N_{\underline{k}\sigma}^* e_{\underline{k}_{nl}\sigma} e^{-i\mathbf{k}_{nl} \cdot \mathbf{h}}) \quad , \quad k_z > 0 \quad . \end{aligned} \quad (4.6)$$

## 7. ASYMPTOTIC EXPANSION

In order to simplify the solution (4.4) for  $\underline{E}_s(\underline{r}, t)$  we consider its value in the radiation zone ( $r \rightarrow \infty$  and  $z > 0$ ). The operators  $b_{\underline{k}\sigma}(t)$  and  $b'_{\underline{k}\sigma}(t)$  are independent of  $\underline{r}$ , so that all  $\underline{r}$ -dependent factors appear in the form  $\exp(i\mathbf{k}_{\alpha} \cdot \underline{r})$ . These exponentials are multiplied by a function of  $\underline{k}_{\alpha}$ , and the result is summed over all values of  $\underline{k}_{\alpha}$ . Therefore, all terms in Eq. (4.4) have the generic form  $V^{-1} \sum g(\underline{k}_{\alpha}) \exp(i\mathbf{k}_{\alpha} \cdot \underline{r})$ . The summation runs either over wave vectors with only positive  $z$ -components or over wave vectors with only negative  $z$ -components. Changing the summation into an integration gives

$$\frac{1}{V} \sum_{\underline{k}_{\alpha}} g(\underline{k}_{\alpha}) e^{i\mathbf{k}_{\alpha} \cdot \underline{r}} = \frac{1}{8\pi^3} \int_0^{\infty} dk_{\alpha} k_{\alpha}^2 \int_{\Omega^{\pm}} d\Omega g(\underline{k}_{\alpha}) e^{i\mathbf{k}_{\alpha} \cdot \underline{r}} \quad , \quad (5.1)$$

where the superscript ( $\pm$ ) on the region of solid angle indicates the sign of the  $z$ -components of the wave vectors  $\underline{k}_{\alpha}$ . Then we can make an asymptotic expansion of the angular integral with the method of stationary phase.<sup>4</sup> The result is

$$\int_{\Omega^+} d\Omega g(\underline{k}_\alpha) e^{i \underline{k}_\alpha \cdot \underline{r}} = \pm 2\pi i \frac{e^{\pm i k_\alpha r}}{k_\alpha r} g(\pm k_\alpha \underline{e}_r) \quad (5.2)$$

where  $\underline{e}_r$  is the radial spherical unit vector which points in the observation direction. This direction will be specified by the spherical angles  $\theta$  and  $\phi$ . The asymptotic expansion effectively filters out the value of  $g(\underline{k}_\alpha)$  for  $\underline{k}_\alpha = \pm k_\alpha \underline{e}_r$ , corresponding to the plane wave  $\exp(i \underline{k}_\alpha \cdot \underline{r})$  which travels into the observation direction.

With Eqs. (5.1) and (5.2), the asymptotic expansion of  $\underline{E}_s(\underline{r}, t)$  is found to be

$$\begin{aligned} \underline{E}_s(\underline{r}, t) = & \frac{-1}{8\pi^2 \epsilon_0 r} \times \sum_{\sigma} \left[ \int_0^\infty dk k e^{-ikr} (\omega e^{-i\omega t} b_{\underline{k}\sigma}(t) \underline{e}_{\underline{k}\sigma})_{\underline{k} = -k \underline{e}_r} \right. \\ & - \int_0^\infty dk_r k_r e^{ik_r r} (\omega e^{-i\omega t} b_{\underline{k}\sigma}(t) R_{\underline{k}\sigma - \underline{k}_r \sigma})_{\underline{k}_r = k_r \underline{e}_r} \\ & + \left\{ \int_0^\infty dk_{pc} k_{pc} e^{-ik_{pc} r} (\omega e^{-i(\omega - 2\bar{\omega})t} b_{\underline{k}\sigma}(t) P_{\underline{k}\sigma - \underline{k}_{pc}\sigma})_{\underline{k}_{pc} = -k_{pc} \underline{e}_r} \right\}^\dagger \\ & - \int_0^\infty dk_t k_t e^{ik_t r} (\omega e^{-i\omega t} b'_{\underline{k}\sigma}(t) T'_{\underline{k}\sigma - \underline{k}_t \sigma})_{\underline{k}_t = k_t \underline{e}_r} \\ & + \left\{ \int_0^\infty dk_{nl} k_{nl} e^{-ik_{nl} r} (\omega e^{-i(\omega - 2\bar{\omega})t} b'_{\underline{k}\sigma}(t) \right. \\ & \left. \left. N'_{\underline{k}\sigma - \underline{k}_{nl}\sigma})_{\underline{k}_{nl} = -k_{nl} \underline{e}_r} \right\}^\dagger \right] + \text{H.c.} \quad (5.3) \end{aligned}$$

## VI. POLARIZATION VECTORS

The polarization vectors  $\underline{e}_{\underline{k}_\alpha \sigma}$ , evaluated for  $\underline{k}_\alpha = \pm k_r \underline{e}_r$ , can be expressed in the spherical unit vectors  $\underline{e}_\theta$  and  $\underline{e}_\phi$ . These calculations are similar (but not identical) to the corresponding calculations in Ref 1 (Sec. IX). Here it is convenient to introduce the notation

$$\underline{e}_{\underline{s}}^+ = +\underline{e}_\phi, \quad \underline{e}_{\underline{p}}^+ = \underline{e}_\theta, \quad (6.1)$$

which will enable us to express the results for s-waves and p-waves in a single formula. We find

$$\begin{aligned} \underline{E}_s(\underline{r}, t) = \frac{1}{8\pi^2 \epsilon_0 r} \sum_{\sigma} \{ \underline{e}_{\underline{s}}^- X_{adv, \sigma} + \underline{e}_{\underline{s}}^+ X_{r, \sigma} + \underline{e}_{\underline{s}}^- X_{pc, \sigma} + \underline{e}_{\underline{s}}^+ X_{t, \sigma} \\ + \underline{e}_{\underline{s}}^- X_{nl, \sigma} \} + \text{H.c.}, \end{aligned} \quad (6.2)$$

with

$$X_{adv, \sigma} = - \int_0^\infty dk \, k \omega \, e^{-ikr - i\omega t} (b_{\underline{k}\sigma}(t))_{\underline{k} = -k \underline{e}_r}, \quad (6.3)$$

$$X_{r, \sigma} = \int_0^\infty dk_r \, k_r \omega \, e^{ik_r r - i\omega t} (b_{\underline{k}\sigma}(t) R_{\underline{k}\sigma})_{\underline{k} = k_r \underline{e}_r}, \quad (6.4)$$

and similar expressions for  $X_{pc, \sigma}$ ,  $X_{t, \sigma}$  and  $X_{nl, \sigma}$ . Apparently, the term proportional to  $X_{adv, \sigma}$  is an advanced (non-causal) contribution to  $\underline{E}_s$ , corresponding to incoming spherical waves. The other four terms are retarded (causal) solutions, and they have the form of outgoing spherical waves.

Next we substitute expressions (4.5) and (4.6) for  $b_{\underline{k}\sigma}(t)$  and  $b'_{\underline{k}\sigma}(t)$ , respectively, into the results for the  $X_{\alpha,\sigma}$ 's. Then we evaluate the polarization vectors in  $b_{\underline{k}\sigma}(t)$  and  $b'_{\underline{k}\sigma}(t)$  at the indicated wave vectors. After lengthy calculations we find

$$\begin{aligned}
 X_{r,\sigma} = & \int_0^\infty dk \, k \omega R_\sigma \times [e^{-i\omega(t-r-r/c)} \int_0^t dt' \, e^{i\omega t'} \, \underline{\epsilon}_\sigma \cdot \underline{\mu}(t')^{(+)} \\
 & + R_\sigma^* e^{-i\omega(t+r-r/c)} \int_0^t dt' \, e^{i\omega t'} \, \underline{\epsilon}_\sigma^+ \cdot \underline{\mu}(t')^{(+)} \\
 & + P_\sigma^* e^{-i\omega(t-r-r/c)} \int_0^t dt' \, e^{i(\omega-2\bar{\omega})t'} \, \underline{\epsilon}_\sigma \cdot \underline{\mu}(t')^{(-)}] , \quad (6.5)
 \end{aligned}$$

and four other similar expressions. Here we introduced the abbreviations

$$\underline{\epsilon}_s^+ = -\underline{e}_\phi , \quad \underline{\epsilon}_p^+ = -\underline{e}_\theta - 2\sin\theta \underline{e}_z , \quad (6.6)$$

$$r = \frac{h}{c} \cos\theta , \quad (6.7)$$

and all Fresnel coefficients, like  $P_\sigma$ ,  $R_\sigma$ , etc., are evaluated at frequency  $\omega$  and an angle of incidence equal to the polar angle ( $\theta$ ) of observation.

## VII. TWO-LEVEL ATOM

So far we have not made any assumption about the atom and its dipole moment  $\underline{\mu}(t)$ . In order to evaluate the expression for  $\underline{E}_s(\underline{r},t)$  further, we assume that only two atomic levels, which might be degenerate, are of

relevance. The level separation is  $\hbar\omega_0$ . In free evolution, the positive and negative frequency part of  $\underline{\mu}(t)$  obey the identity

$$\underline{\mu}(t_2)^{(\pm)} = e^{\mp i\omega_0(t_2-t_1)} \underline{\mu}(t_1)^{(\pm)} \quad (7.1)$$

We shall use this as an approximation in equations of the type (6.5). Then, for instance, Eq. (6.5) can be written as

$$\begin{aligned} X_{r,\sigma} = & \int_0^\infty dk \, k\omega R_\sigma \\ & \times [\epsilon_{\sigma\sim\sigma}^+ \cdot \underline{\mu}(t-r-r/c)^{(+)} e^{-i(\omega-\omega_0)(t-r-r/c)} \int_0^t dt' e^{i(\omega-\omega_0)t'} \\ & + R_{\sigma\sim\sigma}^* e_{\sigma\sim\sigma}^+ \cdot \underline{\mu}(t+r-r/c)^{(+)} e^{-i(\omega-\omega_0)(t+r-r/c)} \int_0^t dt' e^{i(\omega-\omega_0)t'} \\ & + P_{\sigma\sim\sigma}^* \epsilon_{\sigma\sim\sigma}^+ \cdot \underline{\mu}(t-r-r/c)^{(-)} e^{-2i\bar{\omega}(t-r-r/c)} \\ & \times e^{-i(\omega+\omega_0-2\bar{\omega})(t-r-r/c)} \int_0^t dt' e^{i(\omega+\omega_0-2\bar{\omega})t'} ] \quad (7.2) \end{aligned}$$

In every term, both the exponential in the integrand and the exponential in front of the integral have the same frequency. When the distance  $r$  is much larger than a wavelength and the retardation  $r/c$  is much larger than an optical cycle, then the integrals can be approximated by<sup>5</sup>

$$e^{-i\omega_\alpha(t \pm r/c)} \int_0^t dt' e^{i\omega_\alpha t'} = 2\pi \delta(\omega_\alpha) \quad (7.3)$$

In writing out all  $X_{\alpha,\sigma}$  as in Eq. (7.2), it can be shown that every integral can be written as in Eq. (7.3). The only exception is  $X_{adv,\sigma}$ , which contains the integral of Eq. (7.3) with  $r$  replaced by  $-r$ . This makes the right-hand side zero, instead of  $2\pi\delta(\omega_\alpha)$ , as shown in Ref. 5. Therefore,

$$X_{adv,\sigma} = 0, \quad (7.4)$$

as it should. Then we set  $k = \omega/c$  and carry out the integrations over  $\omega$ . This gives, for instance,

$$\begin{aligned} X_{r,\sigma} = & \frac{2\pi\omega_0^2}{c^2} [R_{\sigma-\sigma}^+ \cdot \underline{\mu}(t-r-r/c)^{(+)} + |R_\sigma|^2 e_\sigma^+ \cdot \underline{\mu}(t+r-r/c)^{(+)} \\ & + \bar{R}_\sigma \bar{P}_\sigma^* \cdot \underline{\mu}(t-r-r/c)^{(-)} e^{-2i\bar{\omega}(t-r-r/c)}], \end{aligned} \quad (7.5)$$

where

$$P_\sigma = P_\sigma(\omega_0, \theta), \quad (7.6)$$

$$\bar{P}_\sigma = P_\sigma(2\bar{\omega} - \omega_0, \theta), \quad (7.7)$$

and similarly for other Fresnel coefficients.

### VIII. TOTAL SOURCE FIELD

The total field  $\underline{E}_s(\underline{r}, t)$  has four contributions of the form (7.5), according to Eq. (6.2). We set  $t - r/c \rightarrow t$ , and introduce the polarization vectors (without the  $\pm$  superscripts)

$$\underline{e}_s = \underline{e}_\phi, \quad \underline{e}_p = \underline{e}_\theta, \quad (8.1)$$

$$\underline{\epsilon}_s = \underline{e}_\phi, \quad \underline{\epsilon}_p = -\underline{e}_\theta - 2\sin\theta \underline{e}_z. \quad (8.2)$$

Then we define the parameters

$$\gamma_\sigma^a = |\underline{R}_\sigma|^2 + |\underline{T}'_\sigma|^2 - |\underline{\bar{P}}_\sigma|^2 - |\underline{\bar{N}}'_\sigma|^2, \quad (8.3)$$

$$\gamma_\sigma^i = \underline{\bar{R}}_\sigma \underline{\bar{P}}_\sigma^* - \underline{R}_\sigma \underline{P}_\sigma^* + \underline{\bar{T}}'_\sigma \underline{\bar{N}}'_\sigma^* - \underline{T}'_\sigma \underline{N}'_\sigma^*, \quad (8.4)$$

which will depend on the angle  $\theta$ , in general. When we group together all terms with  $\underline{\mu}(t)^{(+)}$  and all terms with  $\underline{\mu}(t)^{(-)}$ , then the field assumes the remarkably simple form

$$\underline{E}_s(\underline{r}, t) = \frac{\omega_o^2 e^{-i\omega_o t}}{4\pi\epsilon_o c^2 r} \sum_\sigma \underline{e}_\sigma \{ \underline{a}_\sigma^+ \cdot \underline{\mu}(t)^{(+)} + e^{-2i\omega t} \underline{a}_\sigma^- \cdot \underline{\mu}(t)^{(-)} \} + \text{H.c.}, \quad (8.5)$$

in terms of the polarization-like vectors

$$\underline{a}_\sigma^+ = \gamma_\sigma^a \underline{e}_{\sigma-\sigma} + e^{2i\omega_o t} \underline{R}_{\sigma-\sigma} \underline{\epsilon}_\sigma, \quad (8.6)$$

$$\underline{a}_\sigma^- = -\underline{P}_{\sigma-\sigma}^* \underline{e}_\sigma + e^{2i\omega_o t} \gamma_\sigma^i \underline{\epsilon}_{\sigma-\sigma}. \quad (8.7)$$

Alternatively, we can group the terms as

$$\underline{E}_s(\underline{r}, t) = \frac{\omega_o^2 e^{-i\omega_o t}}{4\pi\epsilon_o c^2 r} \{ (\underline{M}_\theta \cdot \underline{e}_\theta) \underline{e}_\theta + (\underline{M}_\phi \cdot \underline{e}_\phi) \underline{e}_\phi \} + \text{H.c.} \quad (8.8)$$

with

$$\begin{aligned} \underline{M}_\theta &= \gamma_{p\mu}^a(t)^{(+)} + e^{2i\omega_o t} R_{p\mu'}(t)^{(+)} \\ &- e^{-2i\omega_o t} (P_{p\mu}^*(t)^{(-)} - e^{2i\omega_o t} \gamma_{p\mu'}^i(t)^{(-)}) \quad (8.9) \end{aligned}$$

$$\begin{aligned} \underline{M}_\phi &= \gamma_{s\mu}^a(t)^{(+)} - e^{2i\omega_o t} R_{s\mu'}(t)^{(+)} \\ &- e^{-2i\omega_o t} (P_{s\mu}^*(t)^{(-)} + e^{2i\omega_o t} \gamma_{s\mu'}^i(t)^{(-)}) \quad (8.10) \end{aligned}$$

The mirror dipole  $\underline{\mu}'$  is defined as

$$\underline{\mu}'(t) = \underline{\mu}_\perp(t) - \underline{\mu}_\parallel(t) \quad (8.11)$$

in terms of the perpendicular and parallel components of  $\underline{\mu}(t)$  with respect to the surface  $z = 0$ . Expressions (8.8) - (8.10) would be identical to our earlier results (12.5) - (12.7) from Ref.1, if the parameters  $\gamma_\sigma^a$  and  $\gamma_\sigma^i$  would be  $\gamma_\sigma^a = 1$  and  $\gamma_\sigma^i = 0$ .

#### IX. SPECIAL CASES

We have not used any of the properties of the Fresnel coefficients in the derivation of the results of the previous section. These coefficients can



be calculated explicitly,<sup>7</sup> but the result is very complicated. Therefore, we consider two limiting cases of practical interest.

#### A. Dielectric Layer

When we turn off the pump beams, then the medium becomes an ordinary dielectric. Therefore,

$$P_{\sigma} = N_{\sigma} = 0 \quad , \quad (9.1)$$

which gives

$$\gamma_{\sigma}^i = 0 \quad . \quad (9.2)$$

The non-zero Fresnel coefficients are related by

$$|T_{\sigma}|^2 + |R_{\sigma}|^2 = 1 \quad , \quad (9.3)$$

so that

$$\gamma_{\sigma}^a = 1 \quad . \quad (9.4)$$

This shows that for a dielectric layer, the results (8.8) - (8.10) are identical to those in Ref.1. Notice that the term which is proportional to  $\mu(t)^{(-)}$  in Eq. (8.5) disappears in this limit.

### B. Transparent PC

When the dielectric constant equals unity, the specular waves vanish. This gives

$$R_{\sigma} = N_{\sigma} = 0 \quad . \quad (9.5)$$

The non-zero Fresnel coefficients are now related by<sup>7</sup>

$$|T_{\sigma}|^2 - |P_{\sigma}|^2 = 1 \quad , \quad (9.6)$$

which holds for any polarization, angle of incidence and frequency. Therefore, we find again

$$\gamma_{\sigma}^a = 1 \quad , \quad \gamma_{\sigma}^i = 0 \quad . \quad (9.7)$$

### X. CONCLUSION

We have derived an expression for the fluorescence radiation field which is emitted by an atomic dipole near the surface of a PC. The starting point was the standard plane-wave expansion of the electric field in terms of annihilation and creation operators. Then we solved the Heisenberg equation of motion for the annihilation operator. This gives rise to two contributions to the electric field: the vacuum field  $\underline{E}_v$  and the source field  $\underline{E}_s$ , which is generated by the dipole. The form of  $\underline{E}_v$  follows trivially from the choice of  $H_r$ , but the form of  $\underline{E}_s$  depends on the choice of interaction Hamiltonian and the structure of  $\underline{E}_v$  (which equals  $\underline{E}(\underline{r}, 0)$ ). We have shown that  $\underline{E}_s$  is identical to the solution of Maxwell's equations, as found previously, provided that

$$\gamma_{\sigma}^a = 1 \quad , \quad \gamma_{\sigma}^i = 0 \quad . \quad (10.1)$$

These parameters do not appear in the solution of Maxwell's equations, which indicates that both approaches are independent indeed. The parameters  $\gamma_{\sigma}^a$  and  $\gamma_{\sigma}^i$  are determined by the Fresnel reflection and transmission coefficients for a plane wave, and they depend on the polarization, frequency and angle of incidence. The general form of the Fresnel coefficients is extremely complicated, which prohibits the verification of Eq. (10.1) for the most general case. We have shown, however, that for a dielectric layer and for a transparent medium the relations in Eq. (10.1) are satisfied, which covers most cases of practical interest.

#### ACKNOWLEDGMENT

This research was supported by the Office of Naval Research and the National Science Foundation under Grant CHE-9016789.

# APPENDIX

Here we summarize the expressions for the various wave vectors in Eq. (2.1) and the phase conventions for the unit polarization vectors. Any wave can be decomposed as

$$\underline{k}_i = \underline{k}_{\parallel} + k_{i,z} \underline{e}_z, \quad (\text{A.1})$$

and any  $\underline{k}_i$  in Eq. (2.1) which corresponds to a given incident wave must have the same parallel component  $\underline{k}_{\parallel}$  with respect to the surface of the medium. The polarization vectors for s-waves and p-waves are chosen as

$$\underline{e}_{k_i s} = \frac{1}{k_{\parallel}} \underline{k}_{\parallel} \times \underline{e}_z, \quad (\text{A.2})$$

$$\underline{e}_{k_i p} = \frac{1}{k_i} \underline{k}_i \times \underline{e}_{k_i s}, \quad (\text{A.3})$$

respectively. An incident wave from  $z > 0$ , and with wave vector  $\underline{k} = k_{\parallel} + k_z \underline{e}_z$  generates a specular (r) wave and a phase-conjugated (pc) wave. The  $z$ -components of their wave vectors are determined by the dispersion relation, and found to be

$$k_{r,z} = -k_z, \quad (\text{A.4})$$

$$k_{pc,z} = -\sqrt{\rho^2 k^2 - k_{\parallel}^2}, \quad (\text{A.5})$$

in terms of

$$\rho = (2\bar{\omega} - \omega)/\omega. \quad (\text{A.6})$$

A wave with wave vector  $\underline{k}$  which is incident on the layer from  $z < 0$ , generates a transmitted (t) wave in  $z > 0$  with wave vector  $\underline{k}$ . In addition, a nonlinear (nl) wave is produced in  $z > 0$ , which has the same frequency shift with respect to the incident wave as the pc-wave.

# REFERENCES

1. H. F. Arnoldus and T. F. George, Phys. Rev. A (1990) submitted.
2. B. H. W. Hendriks and G. Nienhuis, Phys. Rev. A 40, 1892 (1989).
3. B. H. W. Hendriks and G. Nienhuis, J. Mod. Opt. 36, 1285 (1989).
4. M. Born and E. Wolf, Principles of Optics (Pergamon, Oxford, 1980), p. 752.
5. R. Loudon, The Quantum Theory of Light (Clarendon, Oxford, 1983) pp. 53 & 204.
6. H. F. Arnoldus and T. F. George, J. Mod. Opt. 36, 31 (1989).



**UNIVERSITY AT BUFFALO**  
STATE UNIVERSITY OF NEW YORK

Office of the Dean  
Faculty of Natural Sciences  
and Mathematics  
411 Capen Hall  
Buffalo, New York 14260  
(716) 636-2531  
Facsimile: (716) 636-2534  
Bitnet: NSMDEAN@UBVMS

February 11, 1991

Professor Roger W. Anderson  
Department of Chemistry  
University of California  
Santa Cruz, CA 95064

Dear Roger:

Here are some papers to be listed in the upcoming issue #52 of Molecular Dynamics News:

#### **Phase-Conjugated Fluorescence**

*Phys. Rev. A*

Henk F. Arnoldus and Thomas F. George\*

Departments of Chemistry and Physics, SUNY-Buffalo, Buffalo, NY 14260

Fluorescent emission by an atom near a phase conjugator (PC) based on four-wave mixing is studied from first principles, where the Maxwell-Heisenberg equations are solved for the radiation field, and with an asymptotic expansion an expression is derived for the field in the far zone. It is pointed out that an atom in its ground state polarizes the nonlinear medium of the PC, which subsequently can emit spontaneously two photons, and an absorption-emission-absorption process by the atom then produces a fluorescent photon, together with a spontaneous excitation of the atom.

#### **Critical Behavior in Magnetic Superlattices**

*Journal of Magnetism and Magnetic Materials*

T. Hai, Z. Y. Li, D. L. Lin and Thomas F. George\*

Departments of Chemistry and Physics, SUNY-Buffalo, Buffalo, NY 14260

Based on the effective field theory with correlations, the critical behavior of a magnetic superlattice consisting of two different ferromagnets is examined, where a simple cubic Ising model with nearest-neighbor coupling is assumed. It is found that there exists a critical value for the interface exchange coupling above which the interface magnetism appears.

Professor Roger W. Anderson

February 11, 1991

Page 2

**Interface-Phonon-Mediated Magnetopolaronic Effect on Impurity Transition Energies in Quantum Wells**

*Phys. Rev. B*

D. L. Lin, R. Chen and Thomas F. George\*

Departments of Chemistry and Physics, SUNY-Buffalo, Buffalo, NY 14260

Electron interactions with interface phonon modes and strictly-confined bulk phonon modes are considered for the first time to calculate the resonant magnetopolaronic  $1s-2p_z$  transition energy of a hydrogenic impurity in the quantum well of a double heterostructure, where an interaction gap is predicted between the bulk LO and TO frequencies, in contrast to the Fröhlich-type interaction. The results are in good agreement with recent experimental data.

Sincerely,

Thomas F. George



FY90 Abstracts Distribution List for Solid State & Surface Chemistry

Professor John Baldeschwieler  
Department of Chemistry  
California Inst. of Technology  
Pasadena, CA 91125

Professor John Eyler  
Department of Chemistry  
University of Florida  
Gainesville, FL 32611

Dr. Sylvia Johnson  
SRI International  
333 Ravenswood Avenue  
Menlo Park, CA 94025

Professor Paul Barbara  
Department of Chemistry  
University of Minnesota  
Minneapolis, MN 55455-0431

Professor James Garvey  
Department of Chemistry  
State University of New York  
Buffalo, NY 14214

Dr. Zakya Kafafi  
Code 6551  
Naval Research Laboratory  
Washington, DC 20375-5000

Dr. Duncan Brown  
Advanced Technology Materials  
520-D Danury Rd.  
New Milford, CT 06776

Professor Steven George  
Department of Chemistry  
Stanford University  
Stanford, CA 94305

Professor Larry Kesmodel  
Department of Physics  
Indiana University  
Bloomington, IN 47403

Professor Stanley Bruckenstein  
Department of Chemistry  
State University of New York  
Buffalo, NY 14214

Professor Tom George  
Dept. of Chemistry and Physics  
State University of New York  
Buffalo, NY 14260

Professor Max Lagally  
Dept. Metal. & Min. Engineerin  
University of Wisconsin  
Madison, WI 53706

Professor Carolyn Cassady  
Department of Chemistry  
Miami University  
Oxford, OH 45056

Dr. Robert Hamers  
IBM T.J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, N Y 10598

Dr. Stephen Lieberman  
Code 522  
Naval Ocean Systems Center  
San Diego, CA 92152

Professor R.P.H. Chang  
Dept. Matls. Sci. & Engineering  
Northwestern University  
Evanston, IL 60208

Professor Charles Harris  
Department of Chemistry  
University of California  
Berkeley, CA 94720

Professor M.C. Lin  
Department of Chemistry  
Emory University  
Atlanta, GA 30322

Professor Frank DiSalvo  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853

Professor John Hemminger  
Department of Chemistry  
University of California  
Irvine, CA 92717

Professor Fred McLafferty  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853-1301

Dr. James Duncan  
Federal Systems Division  
Eastman Kodak Company  
Rochester, NY 14650-2156

Professor Leonard Interrante  
Department of Chemistry  
Rensselaer Polytechnic Institute  
Troy, NY 12181

Professor Horia Metiu  
Department of Chemistry  
University of California  
Santa Barbara, CA 93106

Professor Arthur Ellis  
Department of Chemistry  
University of Wisconsin  
Madison, WI 53706

Professor Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853

Professor Larry Miller  
Department of Chemistry  
University of Minnesota  
Minneapolis, MN 55455-0431

Professor Mustafa El-Sayed  
Department of Chemistry  
University of California  
Los Angeles, CA 90024

Professor Eugene Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, NC 27514

Professor George Morrison  
Department of Chemistry  
Cornell University  
Ithaca, NY 14853

Professor Daniel Neumark  
Department of Chemistry  
University of California  
Berkeley, CA 94720

Professor Robert Whetten  
Department of Chemistry  
University of California  
Los Angeles, CA 90024

Professor David Ramaker  
Department of Chemistry  
George Washington University  
Washington, DC 20052

Professor R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, CA 90024

Dr. Gary Rubloff  
IBM T.J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, NY 10598

Professor Nicholas Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, PA 16902

Professor Richard Smalley  
Department of Chemistry  
Rice University  
P.O. Box 1892  
Houston, TX 77251

Professor Aaron Wold  
Department of Chemistry  
Brown University  
Providence, RI 02912

Professor Gerald Stringfellow  
Dept. of Matls. Sci.  
& Engineering  
University of Utah  
Salt Lake City, UT 84112

Professor Vicki Wysocki  
Department of Chemistry  
Virginia Commonwealth University  
Richmond, VA 23284-2006

Professor Galen Stucky  
Department of Chemistry  
University of California  
Santa Barbara, CA 93106

Professor John Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburg, PA 15260

Professor H. Tachikawa  
Department of Chemistry  
Jackson State University  
Jackson, MI 39217-0510

Professor William Unertl  
Lab. for Surface Sci.  
& Technology  
University of Maine  
Orono, ME 04469

Dr. Terrell Vanderah  
Code 3854  
Naval Weapons Center  
China Lake, CA 93555

Professor John Weaver  
Dept. of Chem. Eng. & Mat. Sci.  
University of Minnesota  
Minneapolis, MN 55455

Professor Brad Weiner  
Department of Chemistry  
University of Puerto Rico  
Rio Piedras, Puerto Rico 00931